



## Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact [support@jstor.org](mailto:support@jstor.org).

CALCULUS.

400. Proposed by H. S. UHLER, Yale University.

The axis of a prism whose right-section is a regular polygon of apothem  $a$  and of  $n$  sides passes through the center of a sphere of radius  $R$ . Show that, in general, the volume may be expressed by the formula:

$$V = \frac{4}{3} \pi R^3 + \frac{2}{3} a^2 n \left( R^2 - a^2 \sec^2 \frac{\pi}{n} \right)^{1/2} \tan \frac{\pi}{n} + \frac{1}{3} a n (3R^2 - a^2) \sin^{-1} \left[ \frac{2a \left( R^2 - a^2 \sec^2 \frac{\pi}{n} \right)^{1/2} \tan \frac{\pi}{n}}{R^2 - a^2} \right] - \frac{4}{3} \pi R^3 \sin^{-1} \left[ \frac{R \sin \frac{\pi}{n}}{(R^2 - a^2)^{1/2}} \right].$$

Also discuss the two special cases where

$$a = R \cos \frac{\pi}{n}, \text{ and } n = \infty.$$

401. Proposed by LAENAS G. WELD, Pullman, Illinois.

Given a continuum of triangles whose sides are in arithmetical progression, the common difference being  $h$ : (a) The ratio of the mean value of the areas of all the triangles, the mean of whose three sides is not greater than  $\mu$ , to the area of the triangle, the mean of whose three sides is equal to  $\mu$ , is  $(\mu + 2h)/3\mu$ . Indicate the limiting values of this ratio and show that, when it is equal to  $1/2$ , the triangle whose mean side is  $\mu$  is right angled. (b) The ratio of the mean value of the areas of the circles inscribed in all of these triangles, the mean of whose three sides is not greater than  $\mu$ , to that of the circle inscribed in the triangle, the mean of whose three sides is equal to  $\mu$ , has the limiting values  $1/2$  and  $1/3$ . When the triangle whose mean side is  $\mu$  is right angled, the ratio in question is  $4/9$ . (c) Of the circles circumscribed about these triangles the minimum has the radius  $2h$ .

MECHANICS.

319. Proposed by LAENAS G. WELD, Pullman, Illinois.

A hexagonal pencil lies upon the inclined top of a drawing table and is on the point of either rolling or sliding. Find the angle between its direction and the horizontal edge of the table, the coefficient of friction being  $\mu$ .

320. Proposed by CLIFFORD N. MILLS, Brookings, S. Dak.

A heavy uniform chain of length  $l$  is hung over a rough horizontal cylinder of radius  $r$ . Show that one end of the chain will be

$$\frac{2\mu r}{\mu^2 + 1} (e^{\mu\pi} + 1) + l(e^{\mu\pi} - 1)$$

units lower than the other, just when the chain begins to move, the coefficient of friction being  $\mu$ .

NUMBER THEORY.

237. Proposed by NORMAN ANNING, Chilliwack, B. C.

Prove that for three numbers,  $x, y, z$ ,

$$9\Sigma(x - y)^4 = \Sigma(2x - y - z) = 2\Sigma.$$

238. Proposed by CLIFFORD N. MILLS, Brookings, S. Dak.

Determine the rational value of  $x$  that will render  $x^3 + px^2 + qx + r$  a perfect cube. Apply the result to  $x^3 - 8x^2 + 12x - 6$ .